



FISH PASSAGE CENTER

847 NE 19th Avenue, #250, Portland, OR 97232

Phone: (503) 833-3900 Fax: (503) 232-1259

www.fpc.org/

e-mail us at fpcstaff@fpc.org

MEMORANDUM

TO: Gene McPherson

FROM: Brandon R. Chockley

DATE: December 13, 2013

RE: McCall Hatchery Report (1997-2013)

The Fish Passage Center has been marking Chinook from the McCall Hatchery facility over the last several years as part of the Smolt Monitoring Program (SMP) and the Comparative Survival Study (CSS). For purposes of these studies, data are collected on either the juvenile life stage, or both the juvenile and adult life stages. The SMP provides information for in-season management of the hydrosystem and post-season analyses to the federal, state, and tribal fishery agencies. The CSS is a multi-year program that estimates survival rates over different life stages for spring and summer Chinook produced in major hatcheries. We would like to share with you an update of some of the information we developed under these studies for the Chinook used from the McCall Hatchery facility.

Under the Smolt Monitoring Program, information is collected on the timing and migration speed from the hatchery to Lower Granite Dam. In addition, as part of the CSS study, juvenile survival estimates are developed for the hydrosystem between Lower Granite and Bonneville Dams, as well as survival to adulthood of different passage histories.

Table 1 below provides estimates of minimum, median, and maximum travel time from each year's release to Lower Granite Dam. Also provided are estimates of the 95% confidence limits around the estimated median travel time.

Table 1. McCall Hatchery Summer Chinook Travel Times to Lower Granite Dam

Release Date	Migration Year	Travel Time (Days)			Confidence Limits 95%	
		Min	Med	Max	Lower	Upper
20-Mar	1997	9.9	49.4	100.2	49.1	49.7
30-Mar	1998	14.4	36.5	109.7	36.4	36.5
6-Apr	1999	13.8	39.9	129.0	39.7	40.5
5-Apr	2000	11.3	34.1	114.0	34.0	34.2
26-Mar	2001	24.2	48.5	114.8	48.2	48.6
25-Mar	2002	20.2	51.3	82.5	51.0	51.5
31-Mar	2003	12.4	42.0	101.3	41.9	42.1
22-Mar	2004	16.4	43.5	96.1	43.4	43.6
18-Mar	2005	24.4	49.4	93.1	49.4	49.5
21-Mar	2006	17.5	46.1	76.0	45.9	46.3
19-Mar	2007	20.0	47.0	71.8	46.8	47.2
17-Mar	2008	32.9	54.7	100.5	54.5	54.9
24-Mar	2009	17.1	50.1	92.8	49.9	50.4
23-Mar	2010	29.5	49.5	89.4	48.9	50.4
3/22,3/24	2011	14.5	50.2	117.4	50.0	50.3
3/19-3/21	2012	12.9	41.4	86.1	40.9	41.8
3/25-3/28	2013	16.6	44.6	150.1	44.6	44.7

As with last year, we are providing a table that presents the estimated 10%, 50%, and 90% passage dates of McCall yearling summer Chinook juveniles at Lower Granite Dam for each of the years of tagging (Table 2). Also, Figure 1 is provided as an illustration of how the arrival timing of the 2013 smolt release compares to last year's release, as well as the average of the most recent 10-years (2003–2012).

Table 2. Estimated 10%, 50%, and 90% passage dates of McCall Hatchery yearling summer Chinook at Lower Granite Dam.

Migration Year	Release Date(s)	10% Passage Date	50% Passage Date	90% Passage Date
1997	20-Mar	27-Apr	10-May	16-May
1998	30-Mar	28-Apr	6-May	14-May
1999	6-Apr	2-May	16-May	26-May
2000	5-Apr	1-May	9-May	22-May
2001	26-Mar	30-Apr	13-May	17-May
2002	25-Mar	3-May	15-May	21-May
2003	31-Mar	27-Apr	12-May	20-May
2004	22-Mar	27-Apr	4-May	10-May
2005	18-Mar	29-Apr	7-May	12-May
2006	21-Mar	26-Apr	6-May	16-May
2007	19-Mar	25-Apr	5-May	15-May
2008	17-Mar	5-May	11-May	20-May
2009	24-Mar	26-Apr	13-May	21-May
2010	23-Mar	29-Apr	11-May	22-May
2011	3/22, 3/24	6-May	13-May	18-May
2012	3/19-3/21	25-Apr	28-Apr	14-May
2013	3/25-3/26	4-May	9-May	13-May




Figure 1. Cumulative passage timing of McCall Hatchery yearling summer Chinook to Lower Granite Dam.

Figures 2 and 3 are provided below to put into context out-migration conditions that these spring migrants may have experienced over the years. Figure 2 provides the total spring flow volume (April 3–June 20) for the Snake River (as measured at Ice Harbor), along with the average spring spill proportions at each of Lower Granite, Little Goose, Lower Monumental, and Ice Harbor dams, for each migration year. Figure 3 provides the total spring flow volume (April 10–June 30) for the Lower Columbia (as measured at Bonneville), along with the average spring spill proportions at each of McNary, John Day, The Dalles, and Bonneville dams, for each migration year.




Figure 2. Total spring flow volume in the Snake River (at Ice Harbor Dam) and average spill proportion at Lower Granite, Little Goose, Lower Monumental, and Ice Harbor dams. Spring period in the Snake River is April 3–June 20.




Figure 3. Total spring flow volume in the Lower Columbia River (at Bonneville Dam) and average spill proportion at McNary, John Day, The Dalles, and Bonneville dams. Spring period in the Lower Columbia River is April 10–June 30.

Finally, Table 3 below contains estimates calculated in the CSS study of juvenile survival in the hydrosystem between Lower Granite and Bonneville Dams and survival to adulthood of juvenile salmonids in several categories. Those categories are SAR(T), SAR(C₀), and Weighted SAR_{LGR-10-LGR}, where SAR(T) represents smolts transported from Lower Granite, Little Goose, or Lower Monumental Dam, SAR(C₀) represents smolts migrating in river (undetected at Snake

River transportation collector sites), and $SAR_{LGR-10-LGR}$ is a weighted estimate that is obtained by taking the proportion of the total population of smolts (tagged and untagged) at Lower Granite Dam in each study category and multiplying by the respective study category's $SAR_{LGR-10-LGR}$. In effect, the weighted $SAR_{LGR-10-LGR}$ is the estimated SAR for the overall hatchery release (without jacks). The data presented in Table 3 were taken from various chapters and appendices of the 2013 CSS Annual Report, which can be downloaded from <http://www.fpc.org/documents/CSS.html>). Finally, Figure 4 below is a time series of the Weighted $SAR_{LGR-10-LGR}$ over the fourteen years of available data.

Table 3. McCall Hatchery Summer Chinook Survivals from CSS

Release Date	Migration Year	Juvenile			Adult Survival		
		Survival (LGR-BON)	Proportion Transported	T/C Ratio	SAR(T) %	SAR(C ₀) %	Weighted $SAR_{LGR-10-LGR}$
20-Mar	1997	0.43	0.51	1.38	1.51	1.09	1.31
30-Mar	1998	0.56	0.86	1.96	2.69	1.38	2.50
6-Apr	1999	0.52	0.73	1.49	3.59	2.40	3.26
5-Apr	2000	0.61	0.58	1.89	3.88	2.06	3.12
26-Mar	2001	0.27	0.97	31.9	1.24	0.04 ^B	1.20
25-Mar	2002	0.58	0.68	1.44	1.48	1.03	1.34
31-Mar	2003	0.70	0.54	1.47	0.79	0.54	0.68
22-Mar	2004	0.44	0.93	1.59	0.40	0.25	0.39
18-Mar	2005	0.53	0.86	3.02	0.62	0.20 ^C	0.57
21-Mar	2006 ^D	0.60	0.65	1.11	1.15	1.04	1.06
19-Mar	2007 ^D	0.82	0.27	2.09	1.48	0.71	0.90
17-Mar	2008 ^D	0.50	0.52	1.54	1.35	0.88	1.56
24-Mar	2009 ^D	0.57	0.40	2.00	0.76	0.38	0.52
23-Mar	2010 ^D	0.59	0.28	1.37	0.71	0.52	0.58
3/22, 3/24	2011 ^{A, D}	0.57	0.43	1.44	0.30	0.21	0.30
3/19-3/21	2012 ^{D, E}	0.79	0.32	N/A	N/A	N/A	N/A

^A Migration year 2011 is incomplete with Age 2-salt adult returns through 9/15/2013.

^B Assumed $SAR(C_0)$ same as $SAR(C_1)$ for 2001.

^C In-river SAR is combination of groups C_1 and C_0 .

^D Estimates for migration years 2006 through 2011 reflect use of new methodology developed for random pre-assignment of “monitor mode” and “return-to-river mode” operations. See 2010 CSS Annual Report for details.

^E No adult returns to date, only juvenile metrics are available at this time.




Figure 4. Weighted SAR_{LGR-to-LGR} for McCall Hatchery summer Chinook over the last 15 years (1997–2011). Migration year 2011 is incomplete with Age 2-salt adult returns through 9/15/2013.

We hope that the information we have provided regarding the use and application of information from the marked groups over the last several years is of some use to you. If you would like any additional information regarding these releases please feel free to contact us.

Cc: Pete Hassemer, IDF&G
 Bill Tweit, WDFW
 Jay Hesse, Nez Perce
 Tony Nigro, ODFW
 Howard Schaller, USFWS
 FPAC